Patient's name: **Laboratory number:** Paget's disease of nipple: **Diagnostic summary** absent present Non-neoplastic breast changes: Microcalcification: Specimen type: absent present Location: Specify associated pathology: Tumour type: Lesion size Hormone receptor assay status (if performed) Invasive tumour size/s (mm): ER result: ☐ Positive ☐ Negative Total extent of lesion (DCIS + invasive carcinoma)* (mm): Percentage of nuclei stained: % *if greater than invasive tumour size Predominant staining intensity: PR result: Histological grade Positive ■ Negative Percentage of nuclei stained: Mitotic rate (per 10 high power fields): Predominant staining intensity: Mitosis score (1-3): Tubular formation score (1-3): **HER2 status:** Nuclear grade (1-3): Immunohistochemistry score: Histological grade (1-3): ISH (please specify) result: Peritumoural lymphovascular invasion: Sentinel lymph nodes suspicious Metastatic carcinoma present in____ of ____ sentinel lymph present present and extensive nodes examined Involvement of skin/muscle: Involved Size of metastasis Node 1: Yes ■ No absent present Node 2: Yes ■ No Please specify: Node 3: ☐ Yes ■ No DCIS in specimen Node 4: ☐ Yes ■ No Node 5: ☐ Yes ■ No ☐ low intermediate Nuclear grade: Node 6: Yes ■ No Necrosis: Architecture: ___ Extent of extranodal carcinoma (mm): Calcification: _ **Axillary lymph node dissection Resection margins** Metastatic carcinoma present in _ lymph nodes examined Invasive carcinoma at resection margin: Number identified Number involved Yes Level 1: Orientation of involved margin: Distance from nearest Extent of involvement (mm): specified margin(s): Level 2: Level 3: DCIS at resection margin: Apical: Nature of nodal involvement: *Orientation of involved margin:* Distance from nearest Extent of involvement (mm): specified margin(s): focal diffuse LCIS at resection margin: Extent of extranodal carcinoma (mm): No Comments Atypical lobular hyperplasia/lobular carcinoma in situ: Absent Present classical pleomorphic